



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Relationship of heart rate and electrocardiographic time intervals to body mass in horses and ponies**

Schwarzwald, Colin C ; Kedo, Miki ; Birkmann, Katharina ; Hamlin, Robert L

**Abstract:** **OBJECTIVES:** To investigate the relationship of heart rate (HR) and ECG time intervals to body weight (BWT) in healthy horses and ponies. We hypothesized that HR and ECG time intervals are related to BWT. **ANIMALS:** 250 healthy horses of >30 breeds; 5.5 (1-30) y [median (range)]; 479 (46-1018) kg. **METHODS:** Prospective study. Standard base-apex ECGs were recorded while the horses were standing quietly in a box stall. Mean HR over 15 s was calculated and RR interval, PQ interval, QRS duration, and QT interval were measured by a single observer. QT was corrected for differences in heart rate using Fridericia's formula ( $QT(cf) = QT/(3)\sqrt{RR}$ ). The relationship between ECG variables and BWT, age, sex, and RR interval was assessed using multivariate backward stepwise regression analyses. Goodness of fit of the model was improved when using  $\log(BWT)$  compared to BWT. Body weight was overall the strongest predictor of HR and ECG time intervals. Therefore, only  $\log(BWT)$  was included as an independent variable in the final model. The level of significance was  $p = 0.05$ . **RESULTS:** HR ( $R(2) = 0.21$ ) showed a significant negative relationship and PQ ( $R(2) = 0.53$ ), QRS ( $R(2) = 0.23$ ), QT ( $R(2) = 0.14$ ), and QT(cf) ( $R(2) = 0.02$ ) showed significant positive relationships to  $\log(BWT)$ . **CONCLUSIONS:** Small equine breeds undergoing routine ECG recordings have slightly faster heart rates and shorter ECG time intervals compared to larger equine breeds. Although the magnitude of absolute differences may be small, body weight needs to be considered among other factors when comparing HR and ECG time intervals to normal ranges in horses.

DOI: <https://doi.org/10.1016/j.jvc.2012.03.001>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-62355>

Journal Article

Accepted Version

Originally published at:

Schwarzwald, Colin C; Kedo, Miki; Birkmann, Katharina; Hamlin, Robert L (2012). Relationship of heart rate and electrocardiographic time intervals to body mass in horses and ponies. *Journal of Veterinary Cardiology*, 14(2):343-350.

DOI: <https://doi.org/10.1016/j.jvc.2012.03.001>

# Relationship of Heart Rate and Electrocardiographic Time Intervals to Body Mass in Horses and Ponies

Colin C. Schwarzwald<sup>a\*</sup>, PD Dr.med.vet., PhD; Miki Kedo<sup>a</sup>, DVM; Katharina Birkmann<sup>a</sup>, Dr.med.vet.; Robert L. Hamlin<sup>b</sup>, DVM, PhD

<sup>a</sup>Equine Department, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland.

<sup>b</sup>Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State University, 1900 Coffey Road, Columbus, OH 43210, USA.

**Short title:** Heart Rate vs. Body Mass in Horses

**Keywords:** Cardiology, Electrocardiography, Horse, Allometric scaling.

**Previous presentation:** Schwarzwald CC, Kedo M, Hamlin RL: Correlation of heart rate and electrocardiographic time intervals to body mass in horses and ponies. 29<sup>th</sup> annual ACVIM Forum, June 15–18, 2011, Denver, CO, USA. Oral abstract presentation.

**\*Corresponding author:** Colin C. Schwarzwald, PD Dr. med. vet., PhD, Dipl. ACVIM & ECEIM, Equine Department, Vetsuisse Faculty of the University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland. Phone: +41 44 635 86 97. Fax: +41 44 635 89 05. Email: [cschwarzwald@vetclinics.uzh.ch](mailto:cschwarzwald@vetclinics.uzh.ch).

## ABSTRACT

**Objectives** – To investigate the relationship of heart rate (HR) and ECG time intervals to body weight (BWT) in healthy horses and ponies. We hypothesized that HR and ECG time intervals are related to BWT.

**Animals** – 250 healthy horses of >30 breeds; 5.5 (1-30) y [median (range)]; 479 (46-1018) kg.

**Methods** – Prospective study. Standard base-apex ECGs were recorded while the horses were standing quiet in a box stall. Mean HR over 15 sec was calculated and RR interval, PQ interval, QRS duration, and QT interval were measured by a single observer. QT was corrected for differences in heart rate using Fridericia's formula ( $QT_{cf} = QT / \sqrt[3]{RR}$ ). The relationship between ECG variables and BWT, age, sex, and RR interval was assessed using multivariate backward stepwise regression analyses. Goodness of fit of the model was improved when using log(BWT) compared to BWT. Body weight was overall the strongest predictor of HR and ECG time intervals. Therefore, only log(BWT) was included as an independent variable in the final model. The level of significance was  $p=0.05$ .

**Results** – HR ( $R^2=0.21$ ) showed a significant negative relationship and PQ ( $R^2=0.53$ ), QRS ( $R^2=0.23$ ), QT ( $R^2=0.14$ ), and  $QT_{cf}$  ( $R^2=0.02$ ) showed a significant positive relationship to log(BWT).

**Conclusions** – Small equine breeds undergoing routine ECG recordings have slightly faster heart rates and shorter ECG time intervals compared to large equine breeds. Although the magnitude of absolute differences may be small, body weight needs to be considered among other factors when comparing HR and ECG time intervals to normal ranges in horses.

## INTRODUCTION

A negative association between body weight (BWT) and heart rate (HR) has been observed across a wide range of mammalian species.<sup>1-3</sup> Furthermore, it is well known that electrocardiographic (ECG) time intervals vary with heart rate and body weight.<sup>3-6</sup>

However, the association of HR and BWT within a single species containing a wide spectrum of large and small breeds is less clear and still under debate. Contrary to commonly stated clinical experience, the results of two recent studies indicated that there is no significant association between HR and BWT in apparently healthy dogs.<sup>7,8</sup> However, in one of these studies evaluating 24-h ambulatory ECG recordings of 60 healthy dogs between 2 and 80 kg BWT, a strong trend towards a negative association between average HR and BWT ( $p=0.06$ ) was reported.<sup>8</sup>

Within the equine species, small breeds are generally thought to have higher heart rates than large breeds. Historical data compiled from a variety of electrocardiographic and hemodynamic studies<sup>9-27</sup> on a total of 484 horses and ponies suggests that small breeds have a slightly higher resting heart rate compared to large breeds and draft horses, although some of the data were inconsistent (Fig. 1). However, despite the clinical importance of heart rate and the large differences in size among different equine breeds, there is little specific information about normal heart rates and ECG time intervals in horses and ponies in relation to body size.

The goal of this study was therefore to investigate the relationship of HR and ECG time intervals, respectively, to BWT in apparently healthy horses and ponies and to calculate weight-based normal ranges for all variables.

## ANIMALS, MATERIALS AND METHODS

### *Study population*

The study population was chosen prospectively from the hospital population at The Ohio State University Veterinary Teaching Hospital and at the Vetsuisse Faculty of the University of Zurich. All horses were aged  $\geq 1$  y and considered healthy based on history and physical examination, including careful auscultation of heart and lungs. Animals resisting the procedure or showing nervous behavior were excluded from the study.

Two-hundred and fifty horses (96 female, 46 male, 105 male castrated, 3 not recorded) with an age of 5.5 (1-30) years [median (range)] and a body weight of 479 (46-1018) kg were included in the study. The study population included 43 draft horses (Percheron, Belgian, Clydesdale, Shire, Irish, mixed draft); 20 warmblood horses (Oldenburg, Hanoverian, Holstein, Dutch, Polish, Swedish); 36 Thoroughbreds; 30 Standardbreds; 49 Quarter horses, Paint horses and Appaloosas; 40 small breed horses (miniature horse, Pony of the Americas, Icelandic horse, Shetland pony); and 32 other breeds (Arabian, American Saddlebred, Missouri Fox Trotter, Paso Fino, Morgan horse, Tennessee walker, Tinker, Argentinian, Haflinger, mixed breed).

All animals received adequate human care and were treated during the examination according to the ethical guidelines of the universities.

#### *Electrocardiographic recordings*

Physical examinations and ECG recordings were performed by a variety of different observers. During the procedures, all horses were unsedated, standing quiet in a box stall, gently restrained by experienced handlers. ECG leads with alligator clips were placed to obtain a standard base-apex lead. After an accustoming period of approximately 1 minute, a 15 second ECG was recorded and printed on thermosensitive paper using a multiparameter physiologic monitor<sup>a</sup>. Paper speed was set at 50 mm/s (n=246) or at 25 mm/s (n=4).

## *Rhythm analysis and measurements*

Rhythm analysis and measurements of ECG time intervals were performed by a single observer (MK) using the original paper prints. Occurrence of vagal arrhythmias (ie, marked sinus arrhythmia, sino-atrial (SA) block, second-degree atrioventricular (AV) block) or other rhythm events was recorded. Mean heart rate ( $HR_{15''}$ ) during recordings was calculated by multiplying the number of QRS complexes recorded over 15 seconds by factor 4. The following ECG time intervals were subsequently measured: RR interval, PQ interval, QRS duration, and QT interval. The QT interval was corrected for differences in heart rate using Fridericia's correction ( $QT_{cf} = QT / \sqrt[3]{RR}$ ; RR being the RR interval immediately preceding the respective QT interval).

Three consecutive cardiac cycles were measured for each horse and averaged for further analyses. Cycles immediately following an incident of 2<sup>nd</sup> degree AV block were excluded from analyses.

## *Data analysis and statistics*

Graphical presentation, data analyses, and statistics were performed using commercial computer software<sup>b,c,d</sup>. Summary statistics of age and body weight were calculated and reported as median (range).

Multivariate backward stepwise regression analyses were then performed, including HR and ECG time intervals, respectively, as dependent variable and BWT, age, sex, and RR interval (the latter was included for ECG time intervals only) as independent variables. Breed or breed groups were not included in the models because of their strong relationship to body weight (collinearity). Individual temperament and demeanor as well as athletic condition of the horses were not included in the models because of lack of objective measures of these

factors. Validity of the normality assumption was confirmed by assessment of histograms and normal probability plots of the residuals.

Body weight was overall the strongest predictor of HR and ECG time intervals. For all variables except QT interval, inclusion of age, sex, and RR interval in the model did not markedly increase the coefficient of determination (absolute increase of  $R^2 < 0.047$ ; data not shown). For QT (but not QT<sub>cf</sub>), inclusion of RR improved  $R^2$  from 0.144 to 0.331, which was expected because of the well known rate dependence of the QT interval.<sup>9,10,28</sup> Goodness of fit of the model, assessed using the Akaike's Information Criterion, was improved after log transformation of BWT. Based on these preliminary analyses, only log(BWT) was included as an independent variable in the final models. The results of the regression analyses were graphically displayed by scatter plots including the regression line and the 95% confidence band of the regression line. The coefficient of determination ( $R^2$ ) was reported. The normal ranges of HR and ECG time intervals in relation to BWT were graphically displayed by the 95% prediction band. The level of significance was set at  $p = 0.05$ .

## RESULTS

The majority of the horses were in regular sinus rhythm (NSR) during the examination. Fifteen horses (6 %) showed arrhythmias related to high vagal tone, including marked sinus arrhythmia (n=4; 3 small breeds, 1 Quarter yearling; 5–26 years; 80–109 kg), SA block (n=3; 2 Standardbreds, 1 small breed; 2–9 years; 240–471 kg), and second degree AV block (n=8; 3 Thoroughbreds, 3 Quarter horses, 2 others; 5–16 years; 489–723 kg). One horse had an atrial premature complex (APC) and one horse had two ventricular premature complexes (VPC) during the 15-second recording.

The results of the univariate regression analyses are displayed in Fig. 2 and 3. HR<sub>15"</sub> showed a significant negative relationship to log(BWT), whereas RR, PQ, QRS, QT, and QT<sub>cf</sub> showed a significant positive relationship to log(BWT).

## DISCUSSION

The results of the current study indicate that in clinical situations such as during standard ECG recording, resting HR is significantly related to BWT in the equine species. This study therefore is in agreement with general clinical experience and with historical data.<sup>9-27</sup> However, the association between HR and BWT is relatively weak. The absolute difference in mean heart rate is only approximately 4 min<sup>-1</sup> in the weight range between 400 and 700 kg. It increases to 18 min<sup>-1</sup> when considering a larger range between 50 and 1000 kg. Statistically, only about 21% of the variation in HR can be explained by differences in BWT (when analyzed after log transformation of BWT), whereas the remaining 79% must be explained by other factors or inherent variability. This is not surprising, since resting heart rate in healthy animals is known to depend on a variety of other factors, including age, sex, temperament, athletic condition, and possibly breed.<sup>7,29</sup> While a significant influence of age and sex could not be shown in the present study on horses  $\geq 1$  y of age, the effects of temperament and demeanor, athletic condition, and breed were not evaluated.

Some horses might have been slightly stressed because of the hospital environment, the physical restraint, and the data collection procedure. In fact, the prevalence of second degree AV blocks in this study population (8/240, 3.3%) was markedly lower than reported for horses examined at rest (15-23%)<sup>30-32</sup> and for horses undergoing 24-hour Holter ECG monitoring (up to 44%),<sup>33</sup> possibly indicating some degree of vagal withdrawal and/or sympathetic stimulation. Although stress and excitement could have been reduced by using ambulatory Holter ECG or telemetric ECG monitoring, it should be noted that this study more



164 closely reflects the real clinical situation and is therefore representative for the findings  
165 obtained during standard medical procedures. One might argue that ‘cold blooded’ giant  
166 breed horses have a more stolid demeanor and are generally calmer than average size sports  
167 horses, thereby explaining their slightly lower resting heart rate in larger horses. However, the  
168 same might also be true for some small breed horses. Interestingly, marked sinus arrhythmia  
169 and SA block were only observed in horses between 80 and 471 kg BWT and second degree  
170 AV blocks were only observed in horses between 489 and 723 kg BWT. This finding could  
171 indicate that vagal influence is most prominent in horses at a weight range of approximately  
172 80 – 720 kg, but may be less pronounced in very small and very large horses. Alternatively, it  
173 is possible that vagal influence in heavy draft horses preferentially causes slowing of regular  
174 sinus node depolarization rather than SA or AV block. Based on the available information, the  
175 influence of weight- or breed-related differences in temperament cannot be conclusively  
176 determined, all the more considering that individual variation in character independent of  
177 body size is certainly quite large.

178 Athletic condition certainly differs in relation to breed and use of the horses. Small breed  
179 horses (unlike Thoroughbreds, Standardbreds, Quarter horses, and Warmblood horses) are  
180 often not in athletic condition and therefore may have a slightly higher resting heart rate.  
181 However, this may also apply for many giant breed horses, which however, according to the  
182 present study, do have even lower resting heart rates than medium sized horses. Therefore,  
183 population wide differences athletic condition probably cannot explain the weight dependence  
184 of HR found in this study.

185 In mammals, the specific metabolic rate (ie, the metabolic rate per unit mass) decreases  
186 with increasing body size. Hence, energy metabolism is higher in small compared to large  
187 mammals.<sup>34</sup> This has been linked to the inverse relationship between HR and BWT among  
188 mammals, since the increased relative need for oxygen and blood flow in the small animal is

not achieved through a relatively larger heart or stroke volume, but through an increase in heart rate.<sup>35</sup> However, there are some data suggesting that energy metabolism might not be inversely related to body weight within the equine species.<sup>36,37</sup> Hence, differences in specific metabolic rate might not explain the weight dependence of HR in horses and ponies. It has been speculated that differences in sinus node dimensions and the increased coupling of autonomic cells in bigger pacemakers may explain the association of body size to HR in mammals.<sup>38</sup> But the same authors also pointed out that differences in HR between individuals of the same species and age can be quite large, probably as a result of different sets of membrane currents.<sup>38</sup> Yet another group of researchers found that the resting heart rate in mammals might be optimized for pulsatile blood flow in large arteries and therefore is inversely related to body size.<sup>39</sup> In any case, the reasons for the inverse relationship of HR and BWT within the equine species remain largely speculative and it is unknown, whether the explanations that have been formulated for interspecies relationships of HR and BWT in mammals are also applicable to a single mammalian species.

This study also revealed a significant direct relationship between BWT and ECG time intervals. In agreement with previously reported data in horses and ponies,<sup>9-12,40-42</sup> the strongest associations were found between BWT and PQ interval and between BWT and QRS duration. It was not surprising to find that QT interval was not only related to body weight but also to the preceding RR interval.<sup>9,10,28</sup> Fridericia's formula was used exemplarily to eliminate the influence of HR on QT interval in this study. However, rate correction of QT interval is difficult and there is no general consensus on the best formula to use. Many different methods of correction have been proposed for different species and under different conditions.<sup>43-45</sup> In this study, we did not further explore the rate dependence of the QT interval in horses. It is therefore possible that a different method of correction would have been more appropriate or that different methods would have been required for different sex, age, breed or even for

individual horses.<sup>9,44,45</sup> Nonetheless, it was interesting to find that QT<sub>cf</sub> was nearly independent of BWT, suggesting that most of the differences in QT were indeed related to differences in HR, but not to differences in BWT.

As a limitation of this study, general health was determined based on history and physical examination only. No advanced diagnostic methods were used to confirm absence of cardiovascular disease. Therefore, some horses included in the study might have suffered from subclinical, mild cardiovascular disease. In fact, two horses showed atrial and ventricular premature complexes, respectively, possibly indicating cardiac compromise. However, since occasional premature complexes can be observed even in apparently healthy horses,<sup>46-48</sup> this finding is inconclusive. These two horses, that were considered healthy based on medical history and physical examination, were therefore not excluded from the study. Because of the large number of horses in the study population, the error introduced by a few horses possibly suffering from subclinical cardiac disease was considered minor.

## CONCLUSIONS

In conclusion, the present study shows that in clinically healthy horses undergoing routine ECG recordings there is a significant, although weak inverse relationship between HR and BWT. Hence, when observed on a population level, small equine breeds tend to have slightly faster heart rates compared to large breeds. Similarly, there is a significant, but mostly weak direct relationship between BWT and PQ, QRS, QT, and QT<sub>cf</sub> intervals. The strongest association was found between BWT and PQ interval, whereas QT<sub>cf</sub> was nearly independent of BWT. Body weight might have to be considered as one of the determinant factors when interpreting HR measurements and ECG recordings in individual patients under clinical conditions.

239 **FOOTNOTES**

240 <sup>a</sup>Datascope Passport, Datascope Passport, Fumedica AG, Muri, Switzerland.

241 <sup>b</sup>Microsoft Office Excel 2007, Microsoft Corporation, Redmond, WA.

242 <sup>c</sup>GraphPad Prism v5.04 for Windows, GraphPad Software, San Diego, CA.

243 <sup>d</sup>SigmaStat v3.5, SPSS Inc, Chicago, IL.

244

245 **ACKNOWLEDGMENTS**

246       The veterinary anesthesia service of the Veterinary Medical Center of The Ohio State  
247 University and the clinicians of the Equine Department of the Vetsuisse Faculty, University of  
248 Zurich, are gratefully acknowledged for their support with recruitment of horses and  
249 collection of ECG recordings.

250

251 **CONFLICT OF INTEREST**

252       No conflicts of interest to disclose.

## FIGURES

**Figure 1:** Heart rates reported in electrocardiographic and/or hemodynamic studies on a total of 484 horses and ponies, grouped by breed.<sup>9-27</sup> Each dot represents the mean heart rate reported in the respective studies.

**Figure 2:** Results of the univariate regression analyses. *Left:* Semi-logarithmic (lin-log type) scatter plots including the regression line and the 95% confidence band of the regression line. *Right:* Non-logarithmic scatter plots including the regression line and the 95% prediction band. Normal ranges for HR and ECG time intervals for a specific BWT can be estimated based on the prediction band.

$R^2$ , coefficient of determination;  $HR_{15''}$ , mean heart rate calculated over 15 seconds.

**Figure 3:** Results of the univariate regression analyses. *Left:* Semi-logarithmic (lin-log type) scatter plots including the regression line and the 95% confidence band of the regression line. *Right:* Non-logarithmic scatter plots including the regression line and the 95% prediction band. Normal ranges for HR and ECG time intervals for a specific BWT can be estimated based on the prediction band.

$R^2$ , coefficient of determination; RR, RR interval; PQ, PQ interval; QRS, QRS duration; QT, QT interval;  $QT_{cf}$ , rate-corrected QT interval (Fridericia's correction).

## REFERENCES

1. Schmidt-Nielsen K. Circulation. In: Schmidt-Nielsen K, editor. Animal physiology: Adaptation and environment, 5th ed. Cambridge, Cambridge University Press; 1997. p. 91-125.
2. Stahl WR. Scaling of respiratory variables in mammals. J Appl Physiol 1967;22:453-460.
3. Noujaim SF, Lucca E, Munoz V, Persaud D, Berenfeld O, Meijler FL, Jalife J. From mouse to whale: A universal scaling relation for the PR interval of the electrocardiogram of mammals. Circulation 2004;110:2802-2808.
4. Meijler FL, Strackee J, Stokhof AA, Wassenaar C. Scaling of atrioventricular transmission in mammalian species: an evolutionary riddle! J Cardiovasc Electrophysiol 2002;13:826-830.
5. Meijler FL. Atrioventricular conduction versus heart size from mouse to whale. J Am Coll Cardiol 1985;5:363-365.
6. Sawazaki H, Hirose H. Comparative electrocardiographical studies on the conduction time of heart in vertebrates. Jpn J Vet Sci 1974;36:421-426.
7. Ferasin L, Ferasin H, Little CJL. Lack of correlation between canine heart rate and body size in veterinary clinical practice. J Small Anim Pract 2010;51:412-418.
8. Lamb AP, Meurs KM, Hamlin RL. Correlation of heart rate to body weight in apparently normal dogs. J Vet Cardiol 2010;12:107-110.
9. Buss DD, Rwalings CA, Bisgard GE. The normal electrocardiogram of the domestic pony. J Electrocardiol 1975;8:167-172.
10. Lannek N, Rutqvist L. Normal area of variation for the electrocardiogram of horses. Nordisk Vet Med 1951;3:1094-1117.
11. Fregin GF. Electrocardiography. Vet Clin North Am Equine Pract 1985;1:419-432.

- 298 12. Nakamura R, Too K, Matsuhashi A. Studies on applications of electrocardiogram in  
299 horses: I. The lead method and electrocardiogram with special reference to configurations  
300 and measurements of waves in the limb lead. *Jpn J Vet Res* 1955;3:24-32.
- 301 13. Landgren S, Rutqvist L. Electrocardiogram of normal cold blooded horses after work.  
302 *Nord Vet Med* 1953;5:905-914.
- 303 14. Spörri H. [Investigations on the duration of systole and diastole of the heart in domestic  
304 animals and their clinical relevance]. *Schweiz Arch Tierheilkd* 1954;96:593-604.
- 305 15. Matsui K, Sugano S. Relation of intrinsic heart rate and autonomic nervous tone to  
306 resting heart rate in the young and the adult of various domestic animals. *Jpn J Vet Sci*  
307 1989;51:29-34.
- 308 16. Nollet H, Van Loon G, Deprez P, Sustronck B, Muylle E. Use of right ventricular  
309 pressure increase rate to evaluate cardiac contractility in horses. *Am J Vet Res*  
310 1999;60:1508-1512.
- 311 17. Schwarzwald CC, Bonagura JD, Luis-Fuentes V. Effects of diltiazem on hemodynamic  
312 variables and ventricular function in healthy horses. *J Vet Intern Med* 2005;19:703-711.
- 313 18. Grubb TL, Constable PD, Benson GJ, Foreman JH, Olson WO, Thurmon JC, Tranquilli  
314 WJ, Davis LE. Techniques for evaluation of right ventricular relaxation rate in horses and  
315 effects of inhalant anesthetics with and without intravenous administration of calcium  
316 gluconate. *Am J Vet Res* 1999;60:872-879.
- 317 19. Rugh KS, Garner HE, Miramonti JR, Hatfield DG. Left ventricular function and  
318 haemodynamics in ponies during exercise and recovery. *Equine Vet J* 1989;21:39-44.
- 319 20. Orr JA, Bisgard GE, Forster HV, Rawlings CA, Buss D, Will FA. Cardiopulmonary  
320 measurements in nonanesthetized resting normal ponies. *Am J Vet Res* 1975;36:1667-  
321 1670.

- 322 21. Bergsten G. Blood pressure, cardiac output, and blood-gas tension in the horse at rest and  
323 during exercise. *Acta Vet Scand Suppl* 1974;48:1-88.
- 324 22. Durando MM, Reef VB, Birks EK. Right ventricular pressure dynamics during exercise:  
325 relationship to stress echocardiography. *Equine Vet J Suppl* 2002:472-477.
- 326 23. Eberly VE, Gillespie JR, Typler WS. Cardiovascular parameters in the Thoroughbred  
327 horse. *Am J Vet Res* 1964;25:1712-1716.
- 328 24. Steffey EP, Dunlop CI, Farver TB, Woliner MJ, Schultz LJ. Cardiovascular and  
329 respiratory measurements in awake and isoflurane-anesthetized horses. *Am J Vet Res*  
330 1987;48:7-12.
- 331 25. Sexton WL, Erickson HH, Coffman JR. Cardiopulmonary and metabolic responses to  
332 exercise in the Quarter Horse: Effects of training. In: *Proceedings of the 2nd International*  
333 *Conference on Equine Exercise Physiology*, San Diego, CA, 1986.
- 334 26. Goetz TE, Manohar M. Pressures in the right side of the heart and esophagus (pleura) in  
335 ponies during exercise before and after furosemide administration. *Am J Vet Res*  
336 1986;47:270-276.
- 337 27. Hillidge CJ, Lees P. Left ventricular systole in conscious and anesthetized horses. *Am J*  
338 *Vet Res* 1977;38:675-680.
- 339 28. Davey P. How to correct the QT interval for the effects of heart rate in clinical studies. *J*  
340 *Pharmacol Toxicol Methods* 2002;48:3-9.
- 341 29. Hamlin RL, Olsen I, Smith CR, Boggs S. Clinical relevancy of heart rate in the dog. *J*  
342 *Am Vet Med Assoc* 1967;151:60-63.
- 343 30. Buchanan JW. Spontaneous arrhythmias and conduction disturbances in domestic  
344 animals. *Ann N Y Acad Sci* 1965;127:224-238.
- 345 31. Smetzer DL, Smith CR, Senta T. Second-degree atrioventricular block in the horse. *Am J*  
346 *Vet Res* 1969;30:933-946.



- 347 32. Patteson MW, Cripps PJ. A survey of cardiac auscultatory findings in horses. *Equine Vet*  
348 *J* 1993;25:409-415.
- 349 33. Reef VB. The significance of cardiac auscultatory findings in horses: insight into the age-  
350 old dilemma. *Equine Vet J* 1993;25:393-394.
- 351 34. Schmidt-Nielsen K. Metabolic rate and body size. In: Schmidt-Nielsen K, editor. *Scaling:*  
352 *Why is animal size so important.* Cambridge, Cambridge University Press; 1984. p. 56-  
353 74.
- 354 35. Schmidt-Nielsen K. Heart and circulation. In: Schmidt-Nielsen K, editor. *Scaling: Why is*  
355 *animal size so important.* Cambridge, Cambridge University Press; 1984. p. 126-142.
- 356 36. Pagan JD, Hintz HF. Equine energetics. I. Relationship between body weight and energy  
357 requirements in horses. *J Anim Sci* 1986;63:815-821.
- 358 37. Vermorel M, Vernet J, Martin-Rosset W. Digestive and energy utilisation of two diets by  
359 ponies and horses. *Livest Prod Sci* 1997;51:13-19.
- 360 38. Opthof T. The normal range and determinants of the intrinsic heart rate in man.  
361 *Cardiovasc Res* 2000;45:177-184.
- 362 39. Flores J, Corvera Poiré E, del Río JA, López de Haro M. A plausible explanation for  
363 heart rates in mammals. *J Theor Biol* 2010;265:599-603.
- 364 40. Detweiler DK. Electrocardiogram of the Horse. *Fed Proc* 1952;11:34-34.
- 365 41. Deegen E, Reinhard HJ. [Electrocardiographic time patterns in the healthy Shetland  
366 pony]. *Dtsch Tierarztl Wochenschr* 1974;81:257-262.
- 367 42. Reinhard HJ, Zichner M. [Evaluation of telemetrically derived stress electrocardiograms  
368 of the horse using an electronic computer]. *Dtsch Tierarztl Wochenschr* 1970;77:211-  
369 217.
- 370 43. Finley MR, Lillich JD, Gilmour RF, Freeman LC. Structural and functional basis for the  
371 long QT syndrome: Relevance to veterinary patients. *J Vet Intern Med* 2003;17:473-488.

- 372 44. Boyle NG, Weiss JN. Making QT correction simple is complicated. *J Cardiovasc*  
373 *Electrophysiol* 2001;12:421-423.
- 374 45. Goldenberg I, Moss AJ, Zareba W. QT interval: How to measure it and what is “normal”.  
375 *J Cardiovasc Electrophysiol* 2006;17:333-336.
- 376 46. Buhl R, Meldgaard C, Barbesgaard L. Cardiac arrhythmias in clinically healthy  
377 showjumping horses. *Equine Vet J* 2010;42:196-201.
- 378 47. Barbesgaard L, Buhl R, Meldgaard C. Prevalence of exercise-associated arrhythmias in  
379 normal performing dressage horses. *Equine Vet J* 2010;42:202-207.
- 380 48. Ryan N, Marr CM, McGladdery AJ. Survey of cardiac arrhythmias during submaximal  
381 and maximal exercise in Thoroughbred racehorses. *Equine Vet J* 2005;37:265-268.  
382

Figure 1  
[Click here to download high resolution image](#)




Figure 2  
[Click here to download high resolution image](#)





Figure 3  
[Click here to download high resolution image](#)

